

Notes 備忘：

- For contact lenses user, remove contact lenses before applying eye drops.
如配戴隱形眼鏡，用藥前請先把它除下。
- If you are using more than one eye drop or eye ointment, they should be applied 5 to 10 minutes apart.
如同一眼睛需要使用多於一種眼藥水或眼藥膏，兩者需要相隔最少5至10分鐘才能使用。
- Do not share your eye drops with anyone else to avoid transmission of bacteria and delay in treatment.
為防止傳播病菌及避免耽誤病情，切勿與他人共用藥物。

Storage 儲存：

- Store at room temperature unless otherwise suggested on the dispensing label.
請存放於室溫或按照標籤上的指示存放。
- Discard any unused eye drops one month after opening.
若開瓶後一個月仍未將眼藥水用完，應把它棄掉。



For enquiries, please call 如欲查詢，請致電：

Matilda International Hospital - Pharmacy Department
明德國際醫院 — 藥劑部

41 Mount Kellett Road, The Peak, Hong Kong
山頂加列山道41號

t 2849 1531
phar@matilda.org


matilda.org

How to Use Eye Drops 如何使用 眼藥水



To administer a dose 使用方法：

- 1 Wash your hands well before applying eye drops.
首先把雙手徹底清潔。
- 2 Tilt your head backwards and look upwards.
頭部仰後，眼睛向上望。



- 3 Pull the lower eyelid downwards to form a pocket.
用手指輕輕把下眼皮拉下成袋狀。



- 4 Squeeze one drop into the pocket. Do not let the dropper touch against your eye or eyelid to avoid contamination of the dropper.
輕按藥瓶將一滴眼藥水滴入眼袋內。切勿讓瓶嘴接觸眼睛、睫毛或其他東西，以保持瓶嘴清潔。



- 5 Close your eyes and gently press on the inner corner of the eyes for 2 minutes to prevent drops from getting inside the nose and throat.
閉上眼睛並輕按內眼角兩分鐘以防止眼藥水進入鼻及喉嚨。



- 6 Wipe away any excess drops around the eye with a clean tissue.
用清潔的紙巾拭去眼外多餘的藥水。



- 7 Repeat steps 3-6 in the other eye if both eyes are affected.
如雙眼同時需要使用眼藥水，請重覆步驟3至6。
- 8 Tighten the cap after use to prevent contamination.
用後應立刻蓋好藥瓶，以防止藥水受污染。